arXiv:1502.08021v1 [math-ph] 27 Feb 2015

Borzov V.V., Damaskinsky E.V.

The discrete spectrum of Jacobi matrix related to recurrence relations with periodic coefficients[1]

Dedicated to Petr Kulish
in connection with the seventieth anniversary.

1 Introduction

The interest in the study of orthogonal polynomials defined by recurrence relations with periodic and asymptotically periodic coefficients increased after the appearance of the article [1] about the properties of the periodic Toda lattices (see, for example, [2]-[13]). In particular, in work [14] recurrence relation with periodic coefficients are investigated. It was shown that such polynomials can be described by the classical Chebyshev polynomials. The aim of this work is to study the discrete spectrum of the Jacobi matrix, connected with polynomials in this class, i.e. polynomials with periodic coefficients in recurrent relations. As an example, we consider:

a) the case when period N of coefficients in recurrence relations equals to three (as a particular case we consider "parametric" Chebyshev polynomials [18]);

b) the elementary N-symmetrical Chebyshev polynomials ($N = 3, 4, 5$), that was introduced by authors in studying the "composite model of generalized oscillator" [15].

Let us remind some necessary results from [14]. We denote by $\{\varphi_n(x)\}_{n=0}^{\infty}$ the polynomial sequence defined by recurrence relations

$$\varphi_n(x) = (x + a_{n-1})\varphi_{n-1}(x) - b_{n-1}\varphi_{n-2}(x), \quad n \geq 1, \quad \varphi_0(x) = 1, \varphi_{-1}(x) = 0, \tag{1}$$

where the coefficients are periodic with period $N \geq 2$:

$$a_{n+N} = a_n, \qquad b_{n+N} = b_n, \quad n \geq 0. \tag{2}$$

We will use the Chebyshev polynomials of the second kind defined by recurrence relations

$$tU_n(t) = U_{n+1}(t) + U_{n-1}(t), \quad n \geq 0, \quad U_0(t) = 1, U_{-1}(t) = 0. \tag{3}$$

It was proved in [14] that for any $N \geq 1$ the polynomial $\varphi_{N-1}(x)$ divides the polynomial $\varphi_{2N-1}(x)$

$$\varphi_{2N-1}(x) = \varphi_{N-1}(t)P_N(x), \tag{4}$$

---

[1] *Key words*: Orthogonal polynomials, Jacobi matix, recurrent relations.
The work is partially supported by RFBR grant № 15-01-03148-a

where the polynomial $P_N(x)$ defined from equality (4). Besides, recurrence relations

$$\varphi_n(x) = \varphi_{Nm+k}(x) = \varphi_{k+N}(x)U_{m-1}(P_N(x)) - \varphi_k(x)U_{m-2}(P_N(x)). \tag{5}$$

are fulfilled for $n = Nm + k$, $k = \overline{0; (N-1)}$, $m \geq 2$. The Jacobi matrix $J = \left[j_{i,k}^{(N)}\right]_{i,k=0}^{\infty}$ related to recurrence relations (1) one can write in following form

$$j_{i,k}^{(N)} = A\delta_{i+1,k} + B\delta_{i,k} + C\delta_{i-1,k}, \tag{6}$$

where the matrixes

$$A = \begin{pmatrix} 0 & 0 & \dots & 0 \\ 0 & 0 & \dots & 0 \\ \dots & \dots & \dots & \dots \\ 1 & 0 & \dots & 0 \end{pmatrix}, \qquad C = \begin{pmatrix} 0 & 0 & \dots & 0 & b_{N-1} \\ \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & 0 & 0 \\ 0 & 0 & \dots & 0 & 0 \end{pmatrix},$$

$$B = \begin{pmatrix} -a_0 & 1 & 0 & \dots & 0 & 0 \\ b_0 & -a_1 & 1 & \dots & 0 & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & 0 & \dots & 0 & 0 \\ 0 & 0 & 0 & \dots & -a_{N-2} & 1 \\ 0 & 0 & 0 & \dots & b_{N-2} & -a_{N-1} \end{pmatrix}, \tag{7}$$

have the size $N \times N$. Let $X_\mu = (x_1, x_2, \dots, x_n, \dots)^t \in \ell^2$ be an eigenvector for matrix J, corresponding to eigenvalue μ:

$$(J - \mu I)X_\mu = 0. \tag{8}$$

The following necessary and sufficient condition is hold: a solution μ of the equation (8) is an eigenvalue of matrix J if and only if when

$$\|X_\mu\|_2^2 = \sum_{n=1}^{\infty} |x_n|^2 = \sum_{n=0}^{\infty} |\varphi_n(\mu)|^2 < \infty \tag{9}$$

In the next paragraph we will obtain the "critical equation" for Jacobi matrix J . A solution of the equation (9) is called the "critical value" of Jacobi matrix J . The necessary condition for μ to be an eigenvalue of matrix J is that μ be a "critical value" of J .

2 The critical equation for Jacobi matrix J

Let us introduce the notation which was needed in the following:

$$\sigma_{k,k+m} = \sum_{n=k}^{k+m} \varphi_n^2(\mu), \quad S_1 = \sigma_{N,2N-1}, \quad S_2 = \sigma_{0,N-1}, \quad S = S_1 + S_2; \tag{10a}$$

$$S_n^N(\mu) = \sigma_{n,n+2N-1}, \quad D_n^N(\mu) = \sum_{k=n}^{n+N-1} \varphi_k(\mu)\varphi_{k+N}(\mu) \tag{10b}$$

$$\Delta_n^N(\mu) = S_n^N(\mu) - P_N(\mu)D_n^N(\mu), \quad n \geq 0 \tag{10c}$$

It is clear that $S = S_0^N(\mu)$. The following statements are hold:

Lemma 1. *Let the polynomial system* $\{\varphi_n(x)\}_{n=0}^{\infty}$ *is defined by recurrence relations (1) and periodic conditions (2). Then for any* $n \geq 2N$ *the following recurrence relations*

$$\varphi_n(x) = P_N(x)\varphi_{n-N}(x) - \varphi_{n-2N}(x). \tag{11}$$

are fulfilled.

Proof. From relations (3) and (5) we have ($n = Nm + k$)

$$\begin{aligned} P_N(x)\varphi_{n-N}(x) - \varphi_{n-2N}(x) &= \varphi_{k+N}(x)P_N(x)U_{m-2}(P_N(x)) \\ &- \varphi_k(x)P_N(x)U_{m-3}(P_N(x)) - \varphi_{k+N}(x)U_{m-3}(P_N(x)) + \varphi_k(x)U_{m-4}(P_N(x)) = \\ &= \varphi_{k+N}(x)\Big(U_{m-1}(P_N(x)) + U_{m-3}(P_N(x))\Big) - \varphi_k(x)\Big(U_{m-2}(P_N(x)) + U_{m-4}(P_N(x))\Big) \\ &- \varphi_{k+N}(x)U_{m-3}(P_N(x)) + \varphi_k(x)U_{m-4}(P_N(x)) = \\ &= \varphi_{k+N}(x)U_{m-1}(P_N(x)) - \varphi_k(x)U_{m-2}(P_N(x)) = \varphi_{Nm+k}(x) = \varphi_n(x). \end{aligned}$$

□

Lemma 2. *Let the polynomial system* $\{\varphi_n(x)\}_{n=0}^{\infty}$ *is defined by recurrence relations (1) and periodic conditions (2). Besides, let* $\Delta_n = \Delta_n^N(\mu)$ *is defined by the equalities (10b) and (10c). Then for all* $n \geq 0$ *the identity*

$$\Delta_n = \Delta_0 \tag{12}$$

is fulfilled.

Proof. For proof this proposition by induction it is sufficiently to show that for all $n \geq 0$ the equality

$$\Delta_n = \Delta_{n+1}. \tag{13}$$

is fulfilled. Taking into account (10b) and (10c), for proof this relation it is sufficiently to check the equality

$$\begin{aligned} &\varphi_n^2 + \varphi_{n+1}^2 + \ldots + \varphi_{n+2N-1}^2 - P_N\left(\varphi_n\varphi_{n+N} + \varphi_{n+1}\varphi_{n+N+1} + \ldots \varphi_{n+N-1}\varphi_{n+2N-1}\right) = \\ &\quad \varphi_{n+1}^2 + \ldots + \varphi_{n+2N-1}^2 + \varphi_{n+2N}^2 - P_N\left(\varphi_{n+1}\varphi_{n+N+1} + \ldots + \varphi_{n+N-1}\varphi_{n+2N-1} + \varphi_{n+N}\varphi_{n+2N}\right), \end{aligned}$$

This equality is equivalent to the following relation

$$\varphi_n^2 - P_N\varphi_n\varphi_{n+N} = \varphi_{n+2N}^2 - P_N\varphi_{n+N}\varphi_{n+2N}. \tag{14}$$

From (11) it is follow that

$$\varphi_{n+2N}^2 = P_N^2\varphi_{n+N}^2 - 2P_N\varphi_n\varphi_{n+N} + \varphi_n^2. \tag{15}$$

Substituting (15) in (14), we see that the equality (14) is fulfilled. Therefore the equality (13) is also hold. □

Theorem 2.1. *For μ to be an eigenvalue of matrix J defined by (6) and (7) it is necessary that μ be a solution of equation*

$$\Delta_0^N(\mu) = 0. \tag{16}$$

Proof. Let us denote

$$\sigma_{k,k+m} = \sum_{n=k}^{k+m} \varphi_n^2(\mu). \tag{17}$$

It follows from (10), (11) that

$$\sigma_{0,2N-1} = S,$$
$$\sigma_{2N,3N-1} = (P_N\varphi_N - \varphi_0)^2 + \ldots + (P_N\varphi_{2N-1}\varphi_{N-1})^2 = P_N^2 S_1 + S_2 - 2P_N D_0^N(\mu).$$

From (10c) it follows that

$$S - P_N D_0^N = \Delta_0^N \quad \Rightarrow \quad P_N D_0^N = S - \Delta_0^N,$$

so we have

$$\sigma_{2N,3N-1} = 2\Delta_0^N + \left(P_N^2 - 2\right) S_1 - S_2. \tag{18}$$

Analogously, using lemma 2, we obtain

$$\sigma_{3N,4N-1} = 2\Delta_0^N + \left(P_N^2 - 2\right) \sigma_{2N,3N-1} - \sigma_{N,2N-1}. \tag{19}$$

Than we have

$$\sigma_{kN,(k+1)N-1} = 2\Delta_0^N + \left(P_N^2 - 2\right) \sigma_{(k-1)N,kN-1} - \sigma_{(k-2)N,(k-1)N-1}. \tag{20}$$

Summing these equalities in k, we have

$$\begin{aligned}\sigma_{0,nN-1} = \sum_{k=0}^{nN-1} \varphi_k^2 = S + 2\Delta_0^N + \left(P_N^2 - 2\right) S_1 - S_2 \\ + \sum_{k=3}^{n} \left(2\Delta_0^N + \left(P_N^2 - 2\right) \sigma_{(k-1)N,kN-1} - \sigma_{(k-2)N,(k-1)N-1}\right) = \\ = 2(n-1)\Delta_0^N + \left(P_N^2 - 1\right) S_1 + \left(P_N^2 - 2\right)(\sigma_{0,nN-1} - S) - \sigma_{0,nN-1} + S_2 + \sigma_{(n-1)N,nN-1}.\end{aligned}$$

From here we obtain the relation

$$(4 - P_N^2) \sum_{k=0}^{nN-1} \varphi_k^2 = 2(n-1)\Delta_0^N + S_1 + (3 - P_N^2)S_2 + \sigma_{(n-1)N,nN-1}. \tag{21}$$

for finding the quantity $\sigma_{0,nN-1} = \sum_{k=0}^{nN-1} \varphi_k^2$. It is clear that if $\sum_{k=0}^{nN-1} \varphi_k^2(\mu) < \infty$, then

$$\lim_{n\to\infty} \sigma_{(n-1)N,nN-1} = 0. \tag{22}$$

Then it follows from (21) that the series $\sum_{k=0}^{nN-1} \varphi_k^2(\mu)$ is convergent if

$$\Delta_0^N(\mu) = 0.$$

□

Remark 1. All eigenvalues of the matrix J must satisfy the "critical equation" (16).

But it is possible that some critical values of matrix J are not satisfied the necessary and sufficient condition (9), i.e. the corresponding vector X_μ are not belonging to ℓ_2.

Remark 2. Apparently for any $N \geq 2$ the polynomial $\varphi_{N-1}(\mu)$ divides the $\Delta_0^N(\mu)$, i.e.the equality

$$\Delta_0^N(\mu) = \varphi_{N-1}(\mu)\, Q_N(\mu), \tag{23}$$

is true. Then the "critical equation" (16) splits into two equations

$$\varphi_{N-1}(\mu) = 0, \tag{24}$$

and

$$Q_N(\mu) = 0. \tag{25}$$

Apparently that one can obtain a simple condition that a solution μ of equation (24) is an eigenvalue of matrix J. But we are not a success to get a sufficient condition that a solution μ of equation (25) is an eigenvalue of matrix J, which is more simply than the condition (22).

For illustration we consider a few examples. As the first example we consider the matrix J defined by (6), (7) for $N = 3$, $b_0 = b_1 = b_2 = 1$ and for any complex a_0, a_1, a_2.

3 The case $N = 3$. The parametric Chebyshev polynomials

1. Let us consider the generalized Chebyshev polynomials system $\left\{\varphi_n^{(3)}(x)\right\}_{n=0}^{\infty}$ defined by recurrence relations

$$x\varphi_n^{(3)}(x) = \varphi_{n+1}^{(3)}(x) + a_n\varphi_n^{(3)}(x) + \varphi_{n-1}^{(3)}(x), \quad \varphi_0^{(3)}(x) = 1,\ \varphi_{-1}^{(3)}(x) = 0. \tag{26}$$

The coefficients a_n — complex numbers that are fulfilled the periodicity condition (2) with $N = 3$. It is follows from (26) that the first six polynomials are

$$\begin{array}{lll} \varphi_0^{(3)} = 1 & \varphi_1^{(3)} = x - a_0 & \varphi_2^{(3)} = (x - a_1)\varphi_1^{(3)} - 1 \\ \varphi_3^{(3)} = (x - a_2)\varphi_2^{(3)} - \varphi_1^{(3)} & \varphi_4^{(3)} = (x - a_0)\varphi_3^{(3)} - \varphi_2^{(3)} & \varphi_5^{(3)} = \varphi_2^{(3)}(\varphi_3^{(3)} - (x - a_1)), \end{array} \tag{27}$$

and for $n \geq 6$ they can be calculated by formula

$$\varphi_n^{(3)}(x) = P_3(x)\, \varphi_{n-3}^{(3)}(x) - \varphi_{n-6}^{(3)}(x). \tag{28}$$

From (4) and last relation in (27) it follows that

$$P_3(x) = \varphi_3^{(3)}(x) - (x - a_1) = x^3 - (a_0 + a_1 + a_2)x^2 + (a_0a_2 + a_1a_2 + a_0a_1 - 3)x - a_0a_1a_2 + (a_0 + a_1 + a_2). \tag{29}$$

Note that under additional condition $(a_0 + a_1 + a_2) = 0$ the polynomial $P_3(x)$ has more simply form

$$P_3(x) = x^3 + (a_0a_2 + a_1a_2 + a_0a_1 - 3)x - a_0a_1a_2. \tag{30}$$

Let us find the eigenvalues of the matrix $J^{(3)} = \left[j^{(3)}_{j,k}\right]^{\infty}_{j,k=0}$ corresponding to the recurrence relations (26)

$$j^{(3)}_{i\,k} = B_3\delta_{i+1,k} + A_3\delta_{i,k} + B^t_3\delta_{i-1,k}, \tag{31}$$

where

$$A_3 = \begin{pmatrix} a_0 & 1 & 0 \\ 1 & a_1 & 1 \\ 0 & 1 & a_2 \end{pmatrix}, \qquad B_3 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix}. \tag{32}$$

Using the formulas (10a)-(10c), (26) and (27), one can to write the left-hand side of the equation (16) in the following form

$$S^3_0(\mu) - D^3_0(\mu)\,P_3(\mu) = \varphi^{(3)}_2(\mu)\,Q_3(\mu), \tag{33}$$

where

$$Q_3(\mu) = \left(1 + \left(\varphi^{(3)}_1(\mu)\right)^2\right)(\mu - a_1)(\mu - a_2) - 2 + \varphi^{(3)}_2(\mu)\left(1 - \varphi^{(3)}_1(\mu)(\mu - a_2)\right). \tag{34}$$

In result the equation (16) splits into pair of equations

$$\varphi^{(3)}_2(\mu) = \mu^2 - (a_0 + a_1)\mu + a_0a_1 - 1 = 0, \tag{35}$$

$$Q_3(\mu) = 3\mu^2 - 2(a_0 + a_1 + a_2)\mu + (a_0a_2 + a_1a_2 + a_0a_1 - 3) = 0. \tag{36}$$

The roots $\mu_{1,2}$ of the equations (35) have the following form

$$\mu_{1,2} = \mu^{\pm} = \frac{1}{2}(a_0 + a_1) \pm \frac{1}{2}\sqrt{4 + (a_1 - a_0)^2}, \tag{37}$$

and the roots of the equations (36) equal to

$$\mu_{3,4} = \nu^{\pm} = \frac{1}{3}\left((a_0 + a_1 + a_2) \pm \sqrt{a_0^2 + a_1^2 + a_2^2 - a_0a_1 - a_1a_2 - a_0a_2 + 9}\right). \tag{38}$$

In the case $(a_0 + a_1 + a_2) = 0$ the roots $\mu_{3,4}$ are simplified

$$\mu_{3,4} = \pm\sqrt{1 + \frac{a_0^2 + a_1^2 + a_2^2}{6}} \tag{39}$$

It remains to find those critical values μ_k, $(k = 1, 2, 3, 4)$ which are an eigenvalues of Jacobi matrix $J^{(3)}$.

Lemma 3. *For that a root μ_k, $(k = 1, 2)$ (37) of the equation (35) is an eigenvalues of the matrix $J^{(3)}$ (31), (32), it is necessary and sufficient that the following inequality*

$$|\mu_k - a_0| < 1, \quad k = 1, 2, \tag{40}$$

is fulfilled.

Proof. In fact, we have for μ_k, $(k = 1, 2)$ defined by (37) the following relation

$$\sum_{j=0}^{\infty} \left(\varphi_j^{(3)}(\mu_k)\right)^2 = 1 + 2(\mu_k - a_0)^2 \sum_{j=0}^{\infty} (\mu_k - a_0)^{2j}.$$

The series in the right-hand side of this relation is convergent if and only if the following inequality

$$|\mu_k - a_0| < 1, \quad k = 1, 2,$$

is true. $\square$

Unfortunately, for μ_k $(k = 3, 4)$ even in simplest case $(a_0 + a_1 + a_2) = 0$ one cannot find a more simply condition that μ_k is an eigenvalues of the matrix $J^{(3)}$ than the following condition

$$\left[\left(\varphi_{3n}^{(3)}(\mu_k)\right)^2 + \left(\varphi_{3n+1}^{(3)}(\mu_k)\right)^2 + \left(\varphi_{3n+2}^{(3)}(\mu_k)\right)^2\right] \to 0$$

as $n \to \infty$ (but this is the condition (22)).

2. We consider the parametric Chebyshev polynomials introduced in [6] as example to the case when the polynomial $Q_3(\mu)$ has not roots. These polynomials $\{\Psi_n(x; \alpha)\}_{n=0}^{\infty}$ are defined by recurrent relations with coefficients depending on a parameter $\alpha \in [-1, 1]$. These coefficients are

$$a_0(\alpha) = \frac{i\sqrt{3}}{2}(\alpha + 1)(3\alpha - 2), \quad a_1(\alpha) = -i\sqrt{3}\alpha, \quad a_3(\alpha) = -\frac{i\sqrt{3}}{2}(\alpha - 1)(3\alpha - 2), \tag{41a}$$

$$a_{n+3}(\alpha) = a_n(\alpha), \quad n \geq 0, \tag{41b}$$

It is clear, that the following equality

$$a_0(\alpha) + a_1(\alpha) + a_2(\alpha) = 0, \tag{42}$$

is true. From (28)-(32), (41) it is follow that

$$\begin{aligned}
&\Psi_0(x; \alpha) = 1;\\
&\Psi_1(x; \alpha) = x - \frac{i\sqrt{3}}{2}(\alpha + 1)(3\alpha + 2);\\
&\Psi_2(x; \alpha) = x^2 - \frac{}{2}(\alpha - 1)(3\alpha + 2)x + (\frac{3}{2}\alpha(\alpha + 1)(3\alpha - 2) - 1);\\
&\Psi_3(x; \alpha) = x^3 + (1 - \widetilde{\alpha}_1^2)x + i\sqrt{3}\alpha(1 - \widetilde{\alpha}_2^2);\\
&\Psi_4(x; \alpha) = (x - a_0(\alpha))\Psi_3(x; \alpha) - \Psi_2(x; \alpha);\\
&\Psi_5(x; \alpha) = \Psi_2(x; \alpha)P_3(x; \alpha),
\end{aligned} \tag{43}$$

where are used notations

$$\widetilde{\alpha}_1^2 = \frac{27}{4}\alpha^2(1 - \alpha^2), \quad \widetilde{\alpha}_2^2 = \frac{3}{4}(1 - \alpha^2)(9\alpha^2 - 4) \tag{44a}$$

$$P_3(x; \alpha) = x^3 - \widetilde{\alpha}_1^2 x - i\sqrt{3}\alpha\widetilde{\alpha}_2^2. \tag{44b}$$



Figure 1: The support of the continuous spectrum of the Jacobi matrix $J^{(3)}(\alpha)$

For $n \geq 6$ one can calculate the polynomials $\Psi_n(x;\alpha)$ by formulas

$$\Psi_n(x;\alpha) = P_3(x;\alpha)\Psi_{n-3}(x;\alpha) - \Psi_{n-6}(x;\alpha). \tag{45}$$

In the paper [18] we obtain the continuous spectrum of the Jacobi matrix $J^{(3)}(\alpha)$, corresponding to the parametric Chebyshev polynomials. The support of the continuous spectrum is represented on the Fig.1, where are used the notations

$$\lambda_k = \lambda(\alpha)e^{i\frac{2k\pi}{3}}, \qquad \widetilde{\lambda}_k = \lambda(\alpha)e^{i\frac{(2k+3)\pi}{3}}, \quad k = 0,1,2. \tag{46}$$

The number $\lambda(\alpha) \geq 0$ is a positive root of the equation

$$\lambda^3 - \widetilde{\alpha}_1^2\lambda - 2 = 0, \tag{47}$$

where

$$\sqrt[3]{2} \leq \lambda(\alpha) \leq \lambda_{max}, \tag{48}$$

and

$$\lambda_{max} = \sqrt[3]{1+\sqrt{1-\left(\frac{27}{64}\right)^2}} + \sqrt[3]{1-\sqrt{1-\left(\frac{27}{64}\right)^2}}. \tag{49}$$

Now we consider the discrete spectrum of the matrix $J^{(3)}(\alpha)$. From(39) and (41) we find roots of the equation (37)

$$\mu_{1,2}(\alpha) = \frac{i\sqrt{3}}{4}(\alpha-1)(3\alpha+2) \pm \sqrt{1-\frac{3}{16}f^2(\alpha)}, \tag{50}$$

where

$$f(\alpha) = 3\alpha^2 + 3\alpha - 2, \tag{51}$$

and

$$\mu_{1,2}(\alpha) - a_1(\alpha) = -\frac{i\sqrt{3}}{4}f^2(\alpha) \pm \sqrt{1-\frac{3}{16}f^2(\alpha)}. \tag{52}$$

From lemma 3 it follows that μ_k, $k = 1, 2$ is an eigenvalue of the matrix $J^{(3)}(\alpha)$ if and only if the inequality (40) is true. We introduce notations

$$\alpha_1 = -\frac{1}{2} - \sqrt{\frac{1}{4} - \frac{4-2\sqrt{3}}{3\sqrt{3}}},\ \alpha_2 = -\frac{1}{2} + \sqrt{\frac{1}{4} - \frac{4-2\sqrt{3}}{3\sqrt{3}}},\ \alpha_3 = -\frac{1}{2} + \sqrt{\frac{1}{4} + \frac{4+2\sqrt{3}}{3\sqrt{3}}}, \tag{53}$$

and made justification test of the inequality (40) on intervals

$$[-1, \alpha_1], \quad (\alpha_1, \alpha_2), \quad [\alpha_2, \alpha_3], \quad (\alpha_3, 1].$$

As a result we have

a) The numbers $(\mu_{1,2})(\alpha)$ are not eigenvalues of the matrix $J^{(3)}(\alpha)$ as $\alpha \in [-1, \alpha_1)$ or $\alpha \in [\alpha_2, \alpha_3]$.

b) $J^{(3)}(\alpha)$ has only one eigenvalue $\mu_2(\alpha)$ as $\alpha \in (\alpha_1, \alpha_2)$.

c) $J^{(3)}(\alpha)$ has only one eigenvalue $\mu_1(\alpha)$ as $\alpha \in (\alpha_3, 1]$.

Now we consider the solutions $\mu_{3,4}(\alpha)$ of the equation (38), that has in this case the following form

$$Q_3(\mu; \alpha) = 3\mu^2 - \widetilde{\alpha}_1^2 = 0 \tag{54}$$

These solutions are equal to

$$\mu_{3,4}(\alpha) = \pm\frac{\widetilde{\alpha}_1}{\sqrt{3}}. \tag{55}$$

To proof that $\mu_{3,4}(\alpha)$ are not eigenvalues of the matrix $J^{(3)}(\alpha)$ as $\alpha \in [-1, 1]$ it is sufficient to check that the necessary condition (22) is broken. Using the recurrence relations (11) (and notation (10)) it is easy to show that

$$S^N_{3n}(\mu) + S^N_{3(n-1)}(\mu) - S(\mu) - S^N_3(\mu) = P^2_3(\mu; \alpha)\left(\sigma_{3n,3n+2}(\mu) - S_2(\mu)\right).$$

Then

$$\sum_{s=-1}^{1} \sigma_{3(n+s),3(n+s)+2}(\mu) + (1 - P^2_3(\mu; \alpha))\sigma_{3n,3n+2}(\mu) = S_2 + S_1(2 - P^2_3(\mu; \alpha)) + \sigma_{6,6+2}(\mu) = A_0(\alpha). \tag{56}$$

The direct calculation shows that for any $\alpha \in (-1, 0) \cup (0, 1]$ one have $A_0(\alpha) \neq 0$. At the same time the left-hand side of equality (56) tends to zero as $n \to \infty$ if the nesessary condition (22) is fulfilled. From this it follows that $A_0(\alpha) = 0$. Thus $\mu_{3,4}(\alpha)$ are not eigenvalues of the Jacobi matrix $J^{(3)}(\alpha)$ as $\alpha \in (-1, 0) \cup (0, 1]$. From results of [19] it follows that $\mu_{3,4}(0)$ and $\mu_{3,4}(1)$ are not eigenvalues of the Jacobi matrixes $J^{(3)}(0)$ and $J^{(3)}(1)$ too. Therefore, $\mu_{3,4}(\alpha)$ are not eigenvalues of the Jacobi matrix $J^{(3)}(\alpha)$ as $\alpha \in [-1, 1]$.

So the Jacobi matrix $J^{(3)}(\alpha)$ has only one eigenvalue $\mu_1(\alpha)$ for $\alpha \in (\alpha_3, 1]$, and only one eigenvalue $\mu_2(\alpha)$ for $\alpha \in (\alpha_1, \alpha_2)$.

Further, as another example, we consider the Jacobi matrix for the elementary N-symmetric Chebyshev polynomials, which belongs to the type of polynomials under consideration. These polynomials appear in studying of "compound model of generalized oscillator" [15]-[17]. We consider only cases $N = 3, 4, 5$, since it was shown in the paper [15] that such polynomials not exist for $n \geq 6$.

4 Elementary N-symmetric Chebyshev polynomials

Elementary N-symmetric Chebyshev polynomials $\left\{\varphi_n^N(x)\right\}_{n=0}^{\infty}$ [15] are defined by recurrence relations

$$x\varphi_n^N(x) = \varphi_{n+1}^N(x) + a_n\varphi_n^N(x) + \varphi_{n-1}^N(x), \quad \varphi_0^N(x) = 1,\ \varphi_{-1}^N(x) = 0, \tag{57}$$

where coefficients a_n for $N = 3, 4, 5$, given by formulas

$$a_0^{(3)} = i\sqrt{3},\ a_1^{(3)} = i\sqrt{3},\ a_2^{(3)} = 0,\ a_{n+3}^{(3)} = a_n^{(3)},\ n \geq 0; \tag{58a}$$

$$a_0^{(4)} = 2i,\ a_1^{(4)} = 0,\ a_2^{(4)} = -2i,\ a_3^{(4)} = 0,\ a_{n+4}^{(4)} = a_n^{(4)},\ n \geq 0; \tag{58b}$$

$$a_0^{(5)} = a_2^{(5)} = a_3^{(5)} = 0,\ a_1^{(5)} = i\sqrt{5},\ a_4^{(5)} = -i\sqrt{5},\ a_{n+5}^{(5)} = a_n^{(5)},\ n \geq 0. \tag{58c}$$

Using recurrence relations (57), we find first $2N$ $(N = 3, 4, 5)$polynomials

$$\begin{array}{lll} \varphi_0^{(3)}(x) = 1, & \varphi_1^{(3)}(x) = x - i\sqrt{3}, & \varphi_2^{(3)}(x) = x^2 + 2, \\ \varphi_3^{(3)}(x) = x\varphi_2^{(3)}(x) - \varphi_1^{(3)}(x), & \varphi_4^{(3)}(x) = x^3\varphi_1^{(3)}(x) + 1, & \varphi_5^{(3)}(x) = x^3\varphi_2^{(3)}(x); \end{array} \tag{59}$$

$$\begin{array}{ll} \varphi_0^{(4)}(x) = 1, & \varphi_1^{(4)}(x) = x - 2i, \\ \varphi_2^{(4)}(x) = x^2 - 2ix - 1, & \varphi_3^{(4)}(x) = x^3 + 2x, \\ \varphi_4^{(4)}(x) = x^4 + x^2 + 2ix + 1, & \varphi_5^{(4)}(x) = x^5 - 2ix^4 + 3x - 2i, \\ \varphi_6^{(4)}(x) = x^6 - 2ix^5 - x^4 + 2x^2 - 4ix - 1, & \varphi_7^{(4)}(x) = (x^4 + 2)\varphi_3^{(4)}(x); \end{array} \tag{60}$$

$$\begin{array}{ll} \varphi_0^{(5)}(x) = 1, & \varphi_1^{(5)}(x) = x, \\ \varphi_2^{(5)}(x) = x^2 - i\sqrt{5}x - 1, & \varphi_3^{(5)}(x) = x^3 - i\sqrt{5}x^2 - 2x, \\ \varphi_4^{(5)}(x) = x^4 - i\sqrt{5}x^3 - 3x^2 + i\sqrt{5}x + 1, & \varphi_5^{(5)}(x) = x^5 + x^3 - i\sqrt{5}x^2 - 2x + i\sqrt{5}, \\ \varphi_6^{(5)}(x) = x^6 + x^2 - 1, & \varphi_7^{(5)}(x) = x^7 - i\sqrt{5}x^6 - x^5 + x, \\ \varphi_8^{(5)}(x) = x^8 - i\sqrt{5}x^7 - 2x^6 + 1, & \varphi_9^{(5)}(x) = x^5\varphi_4^{(5)}(x). \end{array} \tag{61}$$

From here and relation (4) we get the following expression for $P_N(x)$

$$P_3(x) = x^3, \quad P_4(x) = x^4 + 2, \quad P_5(x) = x^5. \tag{62}$$

In view of (62), the relation (5) takes the following form $(k = 0, 1, 2,\ m \geq 2)$

$$\varphi_{3m+k}^{(3)}(x) = \varphi_{k+3}^{(3)}(x)U_{m-1}(x^3) - \varphi_k^{(3)}(x)U_{m-2}(x^3); \tag{63a}$$

$$\varphi_{4m+k}^{(4)}(x) = \varphi_{k+4}^{(4)}(x)U_{m-1}(x^4 + 2) - \varphi_k^{(4)}(x)U_{m-2}(x^4 + 2); \tag{63b}$$

$$\varphi_{5m+k}^{(5)}(x) = \varphi_{k+5}^{(5)}(x)U_{m-1}(x^5) - \varphi_k^{(5)}(x)U_{m-2}(x^5). \tag{63c}$$

The Jacobi matrix $J^{(N)} = \left[j^{(N)}_{j,k}\right]^{\infty}_{j,k=0}$, $(N = 3,4,5)$ corresponding to the relations (57), has the form

$$J^{(N)} = B_N\delta_{i+1,k} + A_N\delta_{i,k} + B^t_N\delta_{i-1,k}, \tag{64}$$

where

$$A_3 = \begin{pmatrix} i\sqrt{3} & 1 & 0 \\ 1 & -i\sqrt{3} & 1 \\ 0 & 1 & 0 \end{pmatrix}, \qquad B_3 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix}; \tag{65a}$$

$$A_4 = \begin{pmatrix} 2i & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 \\ 0 & 1 & -2i & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix}, \qquad B_4 = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix}; \tag{65b}$$

$$A_4 = \begin{pmatrix} 0 & 1 & 0 & 0 & 0 \\ 1 & i\sqrt{5} & 1 & 0 & 0 \\ 0 & 1 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 1 & -i\sqrt{5} \end{pmatrix}, \qquad B_4 = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 \end{pmatrix}; \tag{65c}$$

Now we turn to evaluation of eigenvalues of the matrix $J^{(N)}$, $(N = 3,4,5)$, using the critical equation (16).

A. Discrete spectrum of Jacobi matrix $J^{(3)}$

The Jacobi matrix $J^{(3)}$ is defined by equalities (64) and (65a). From (33), (59), (62), (36), (10a), taking into account that

$$a^{(3)}_0 + a^{(3)}_1 + a^{(3)}_2 = 0, \qquad a^{(3)}_0 a^{(3)}_1 + a^{(3)}_1 a^{(3)}_2 + a^{(3)}_0 a^{(3)}_2 = 3,$$

we obtain that left-hand side $\Delta^{(3)}_0(\mu)$ of equation (16) has the following form

$$\Delta^{(3)}_0(\mu) = S^{(3)}_0(\mu) - D^{(3)}_0(\mu)\, P_3(\mu) = 3\mu^2(\mu^2+2).$$

Then the equation (16) for $N = 3$ looks as

$$\mu^2(\mu^2+2) = 0.$$

The solutions of this equation are equal to

$$\mu_1 = i\sqrt{2}, \quad \mu_2 = -i\sqrt{2}, \quad \mu_{3,4} = 0. \tag{66}$$

Using the lemma 4, we have

$$|\mu_1 - a_0| = |i\sqrt{2} - i\sqrt{3}| < 1.$$

It means that μ_1 is an eigenvalue of the matrix $J^{(3)}$. Further,

$$|\mu_2 - a_0| = |-i\sqrt{2} - i\sqrt{3}| > 1,$$

i.e. μ_2 is not an eigenvalue of the matrix $J^{(3)}$. Now we calculate the components of vector $X_{\mu_3} = X_{\mu_4}$ for $\mu_{3,4}$. They are equal to

$$x_1 = 1,\, x_2 = -i\sqrt{3},\, x_3 = 2,\, x_4 = i\sqrt{3},\, x_5 = 1,\, x_6 = 0, \quad x_{k+6} = -x_k, \quad \text{при} \quad k \geq 0. \tag{67}$$

Taking into account that $\varphi^{(3)}_{k-1}(\mu) = x_k$, for $k = 3, 4$, we have

$$\left[\left(\varphi^{(3)}_{3n}(\mu_k)\right)^2 + \left(\varphi^{(3)}_{3n+1}(\mu_k)\right)^2 + \left(\varphi^{(3)}_{3n+2}(\mu_k)\right)^2\right] \nrightarrow 0, \quad \text{при} \quad n \to \infty,$$

i.e. the condition (22) is not realized. Therefore $X_{\mu_k} \notin \ell^2$, it means that $\mu_{3,4}$ are not eigenvalues of the matrix $J^{(3)}$.

B. Discrete spectrum of Jacobi matrix $J^{(4)}$

The Jacobi matrix $J^{(4)}$ is defined by equalities (64) and (65b). Using (33), (60), (62), (36), (10b), we rewrite the equation (16) for $N = 4$ in the form

$$\mu^4(\mu^2 + 2) = 0. \tag{68}$$

The solutions of this equation are equal to

$$\mu_1 = i\sqrt{2}, \quad \mu_2 = -i\sqrt{2}, \quad \mu_{3,4,5,6} = 0.$$

In the first place we consider zero solutions of this equation. The components of the vector X_{μ_k} (for $k = 3, 4, 5, 6$) are equal to

$$x_1 = 1,\, x_2 = -\frac{i}{2},\, x_3 = -1,\, x_4 = -\frac{3i}{2}, \quad x_{k+4} = x_k, \quad \text{при} \quad k \geq 0.$$

Then partial sum of the series

$$\sum_{n=0}^{\infty} \left(\varphi^{(4)}_n(\mu_k)\right)^2 = \sum_{n=1}^{\infty} x_k^2$$

equal to

$$S_1 = 1,\, s_2 = 1 - \frac{i}{2},\, S_3 = -\frac{i}{2},\, S_4 = -2i, \quad S_{n+4} = S_n - 2i, \quad \text{при} \quad n \geq 1.$$

Thus the sequence of partial sums of series $\sum_{n=1}^{\infty} x_k^2$ has not limit point as $n \to \infty$, i.e. the series is divirgent. Hence, $\mu_{3,4,5,6}$ are not eigenvalues of the matrix $J^{(4)}$. The components of vector $X_{\mu_1} = (x_1, x_2, \ldots)^t$, corresponding to critical value μ_1, are equal to

$$x_1 = 1,\, x_2 = i(\sqrt{2} - 2),\, x_3 = (2\sqrt{2} - 3),\, x_4 = 0, \quad x_{k+4} = (3 - 2\sqrt{2})x_k, \quad \text{при} \quad k \geq 1.$$

Because

$$\|X_{\mu_1}\|^2 = \sqrt{2},$$

then critical value $\mu_1 = i\sqrt{2}$ is eigenvalue of the Jacobi matrix $J^{(4)}$. The normalized eigenvector $Y_{\mu_1} = (y_1, y_2, \ldots)^t$ has the following components

$$y_{4k+1} = \frac{1}{\sqrt[4]{2}}\,(3-2\sqrt{2})^k,\ y_{4k+2} = \frac{i(\sqrt{2}-2)}{\sqrt[4]{2}}\,(3-2\sqrt{2})^k,\ y_{4k+3} = \frac{(2\sqrt{2}-3)}{\sqrt[4]{2}}\,(3-2\sqrt{2})^k,\ y_{4k+4} = 0,$$

with $k \geq 0$. Finally, the components of vector $X_{\mu_2} = (x_1, x_2, \ldots)^t$, corresponding to critical value $\mu_2 = -i\sqrt{2}$, are equal to

$$x_1 = 1,\ x_2 = -i(2+\sqrt{2}),\ x_3 = -(3+2\sqrt{2}),\ x_4 = 0, \quad x_{k+4} = (3+2\sqrt{2})x_k, \quad \text{при} \quad k \geq 1.$$

Consequently,

$$\|X_{\mu_2}\|^2 = \infty$$

i.e. critical value $\mu_2 = -i\sqrt{2}$ is not eigenvalue of the Jacobi matrix $J^{(4)}$.

C. Discrete spectrum of Jacobi matrix $J^{(5)}$

We consider now the Jacobi matrix $J^{(5)}$ that is defined by equalities (64) and (65c). Using (33), (61), (62), (36), (10c), we rewrite the equation (16) for $N = 5$ in the form

$$\mu^4(\mu^4 - i\sqrt{5}\mu^3 - 3\mu^2 + i\sqrt{5}\mu + 1) = 0. \tag{69}$$

The solutions of this equation are equal to

$$\begin{gathered} \mu_{1,2} = \frac{1}{4}\left[\pm\sqrt{10-2\sqrt{5}} + i(1+\sqrt{5})\right], \\ \mu_{3,4} = \frac{1}{4}\left[\pm\sqrt{10+2\sqrt{5}} + i(-1+\sqrt{5})\right], \\ \mu_{5,6,7,8} = 0. \end{gathered} \tag{70}$$

From the same arguments as given above, we see that for $k = \overline{5;8}$ the vector $X_{\mu_k} \notin \ell^2$, i.e. corresponding critical values μ_k are not eigenvalues of the Jacobi matrix $J^{(5)}$.

For the critical value $\mu_1 = \frac{1}{4}\left[\sqrt{10-2\sqrt{5}} + i(1+\sqrt{5})\right]$ the squared components of vector $X_{\mu_1} = (x_1, x_2, \ldots)^t$ are equal to

$$\begin{gathered} (x_1)^2 = 1, \quad (x_2)^2 = \frac{1-\sqrt{5}}{4} + \frac{i(1+\sqrt{5})}{8}\sqrt{10-2\sqrt{5}}, \\ (x_3)^2 = \frac{\sqrt{5}-2}{2} + \frac{i(1-\sqrt{5})}{8}\sqrt{10-2\sqrt{5}}, \quad (x_4)^2 = \frac{\sqrt{5}-3}{2}, \\ (x_5)^2 = 0, \qquad (x_{k+5})^2 = \frac{\sqrt{5}-3}{2}\,(x_k)^2, \quad k \geq 1, \end{gathered}$$

Then $\|X_{\mu_1}\|^2 = 2\sqrt{5}$ and consequently, μ_1 is an eigenvalue of the Jacobi matrix $J^{(5)}$.

Similarly, for critical value $\mu_2 = \frac{1}{4}\left[-\sqrt{10-2\sqrt{5}} + i(1+\sqrt{5})\right]$ the squared components of vector X_{μ_2} are equal to

$$(x_1)^2 = 1, \quad (x_2)^2 = \frac{1-\sqrt{5}}{4} - \frac{i(1+\sqrt{5})}{8}\sqrt{10-2\sqrt{5}},$$

$$(x_3)^2 = \frac{\sqrt{5}-2}{2} + \frac{i(\sqrt{5}-1)}{8}\sqrt{10-2\sqrt{5}}, \quad (x_4)^2 = \frac{\sqrt{5}-3}{2},$$

$$(x_5)^2 = 0, \qquad (x_{k+5})^2 = \frac{\sqrt{5}-3}{2}\,(x_k)^2, \quad k \geq 1.$$

Therefore, $\|X_{\mu_2}\|^2 = 2\sqrt{5}$ and μ_2 is an eigenvalue of the Jacobi matrix $J^{(5)}$.

For critical value $\mu_3 = \frac{1}{4}\left[\sqrt{10+2\sqrt{5}} + i(\sqrt{5}-1)\right]$ the squared components of vector X_{μ_3} are equal to

$$(x_1)^2 = 1, \quad (x_2)^2 = \frac{1+\sqrt{5}}{4} + \frac{i(\sqrt{5}-1)}{8}\sqrt{10+2\sqrt{5}},$$

$$(x_3)^2 = -\frac{2+\sqrt{5}}{2} - \frac{i(1+\sqrt{5})}{8}\sqrt{10+2\sqrt{5}}, \quad (x_4)^2 = -\frac{\sqrt{5}+3}{2},$$

$$(x_5)^2 = 0, \qquad (x_{k+5})^2 = -\frac{3+\sqrt{5}}{2}\,(x_k)^2, \quad k \geq 1.$$

From these relations follow that $X_{\mu_3} \notin \ell^2$, and corresponding critical value μ_3 is not eigenvalue of the Jacobi matrix $J^{(5)}$.

Finally, for critical value $\mu_4 = -\frac{1}{4}\left[\sqrt{10+2\sqrt{5}} - i(\sqrt{5}-1)\right]$ the squared components of vector X_{μ_4} are equal to

$$(x_1)^2 = 1, \quad (x_2)^2 = \frac{1+\sqrt{5}}{4} - \frac{i(\sqrt{5}-1)}{8}\sqrt{10+2\sqrt{5}},$$

$$(x_3)^2 = -\frac{2+\sqrt{5}}{2} + \frac{i(1+\sqrt{5})}{8}\sqrt{10+2\sqrt{5}}, \quad (x_4)^2 = -\frac{\sqrt{5}+3}{2},$$

$$(x_5)^2 = 0, \qquad (x_{k+5})^2 = -\frac{3+\sqrt{5}}{2}\,(x_k)^2, \quad k \geq 1.$$

As above, from these relations it follows that $X_{\mu_4} \notin \ell^2$ and corresponding critical value μ_4 is not eigenvalue of the Jacobi matrix $J^{(5)}$.

References

[1] M. Kac, P. Van Moerbeke, *On some periodic Toda lattices*. Proc. Nat. Acad. Sci. U.S.A. (2nd ed.), **72**, 1627–1629 (1975).

[2] Аптекарев А. И. *Асимптотические свойства многочленов, ортогональных на системе контуров, и периодические движения цепочек Тода.* Мат. Сб. **125(167)**, № 2(10), 231–258 (1984). (Math. USSR-Sb., 53, pp. 233–260 (1986)).

[3] W. Van Assche, *Christoffel functions and Tura'n determinants on several intervals.* J. Comput. and Appl. Math. **48**:1–2, 207–223 (1993).

[4] Д. Барриос, Г. Лопес, Э. Торрано, *Полиномы, порожденные трехчленным рекуррентным соотношением с асимптотически периодическими комплексными коэффициентами.* Мат. Сб. **186**:5, 3–34 (1995) (Mat. Sb., **186**:5, 3–34 (1995)).

[5] J. Bazargan, I. Egorova, *Jacobi operator with step-like asymptotically periodic coefficients.* Mat. Fiz. Anal. Geom., **10**:3, 425–442 (2003).

[6] J. Geronimo, W. Van Assche, *Orthogonal polynomials with asymptotically periodic recurrence coefficients.* J. Approx. Theory **46**, 251–283 (1986).

[7] J. Gilewicz, E. Leopold, *Zeros of polynomials and recurrence relations with periodic coefficients.* J. Comput. Appl. Math. **107**:2, 241–255 (1999)

[8] C.C. Grosjean, *The measure induced by orthogonal polynomials satisfying a recursion formula with either constant or periodic coefficients.* Part I: *Constant coefficients.* Med. Konink. Acad. Wetensch. Belgie, **48**:3, 39–60 (1986).

[9] P Van Moerbeke, *The spectrum of Jacobi matrices.* Invent. Math. **37**:1, 45-81 (1976).

[10] F. Peherstorfer, *On Bernstein-Szego orthogonal polynomials on several intervals.* II. *Orthogonal polynomials with periodic recurrence coefficients.* J. Approx. Theory **64**:2, 123–161 (1991).

[11] F. Peherstorfer, R. Steinbauer, *Orthogonal polynomials on arcs of the unit circle.* II. *Orthogonal polynomials with periodic reflection coefficients.* J. Approx. Theory **87**:1, 60–102 (1996).

[12] F. Peherstorfer, R. Steinbauer, *Asymptotic Behaviour of Orthogonal Polynomials on the Unit Circle with Asymptotically Periodic Reflection Coefficients.* J. Approx. Theory **88**:3, 316–353 (1997).

[13] A. Almendral Va'zquez, *The Spectrum of a Periodic Complex Jacobi Matrix Revisited.* J. Approx. Theory **105**:2, 344–351 (2000).

[14] B. Beckermann, J. Gilewicz, E. Leopold, *Recurrence relation with periodic coefficients and Chebyshev polynomials*, Applicationes Mathematicae **23**, 319-323 (1995).

[15] В.В. Борзов, Е.В. Дамаскинский, *N-симметричные полиномы Чебышева в составной модели обобщенного осциллятора*, ТМФ **129**, №2, 229-240 (2011).

[16] В.В. Борзов, Е.В. Дамаскинский, *Составная модель обобщенного осциллятора. I* , ЗНС ПОМИ **374**, 58-81 (2010).

[17] V.V. Borzov, E.V. Damaskinsky, *Connection between representations of nonstandard and standard Chebyshev oscillators*, Day on Diffraction 2010 28-34.

[18] V.V. Borzov, E.V. Damaskinsky, *The differential equation for generalized parametric Chebyshev polynomials*, Days on Diffraction 2012.

[19] В.В. Борзов, Е.В. Дамаскинский, *Дифференциальные уравнения для простейших 3-симметричных полиномов Чебышева.* Записки Научных Семинаров ПОМИ, т.398,64-86 (2012).